

Muon Scattering at 219 GeV and the Proton Structure Functions

B. A. Gordon, T. W. Quirk, E. L. Anderson, N. E. Booth, W. R. Francis, R. G. Hicks,
 W. W. Kinnison, T. B. W. Kirk, W. A. Loomis, H. S. Matis, L. W. Mo,
 L. C. Myriantopoulos, F. M. Pipkin, J. E. Hudfoot, A. L. Sessoms,
 W. D. Shambroom, A. Skuja, M. A. Staton, C. Tao,
 W. S. C. Williams, Richard Wilson,
 and S. C. Wright

*Enrico Fermi Institute and University of Chicago, Chicago, Illinois 60637, and Fermilab, Batavia, Illinois 60510
 and High Energy Physics Laboratory and Department of Physics, Harvard University, Cambridge,
 Massachusetts 02138, and Department of Physics, The University of Illinois at
 Urbana-Champaign, Urbana, Illinois 61801, and Department of Physics and
 Astronomy, University of Maryland, College Park, Maryland 20742, and
 Physics Department, Michigan State University, East Lansing,
 Michigan 48824, and Department of Nuclear Physics, The
 University of Oxford, Oxford OX1 3RH, England, and
 Physics Department, Virginia Polytechnic Institute
 and State University, Blacksburg, Virginia 24061*

(Received 21 June 1978)

Results on the proton structure function, F_2 , are presented for $0.3 < q^2 < 80.0 \text{ GeV}^2$ and $10 < \nu < 200 \text{ GeV}$. The results support the conclusions of earlier work at 97 and 147 GeV that scaling is violated. A new value for $R = \sigma_S/\sigma_T = 0.44 \pm 0.25$ has been obtained using all the Fermilab proton measurements.

In this Letter we report the most recent measurements at Fermilab of the cross section for scattering of 219-GeV muons from a hydrogen target. The data were taken using the muon-scattering facility, details of which have been published previously.^{1,2} The results are based on a total of 7.5×10^{10} incident muons yielding 2.0×10^4 useful events in the kinematic range $0.3 < q^2 < 80 \text{ GeV}^2$ and $10 < \nu < 200 \text{ GeV}$, where q^2 is the square of the four-momentum transfer of the muon and ν its laboratory energy loss. There were 1.5×10^4 events with $q^2 > 1.0 \text{ GeV}^2$.

The cross section for muon inclusive scattering, in the one-photon-exchange approximation, is related to the structure function F_2 by

$$\frac{d^2\sigma}{dq^2 d\nu} = \frac{2\pi\alpha^2}{p^2 q^4} \frac{F_2(q^2, \nu)}{\nu} \left[2EE' - \frac{q^2}{2} + \frac{(q^2 - 2m_\mu^2)(1 + \nu^2/q^2)}{1 + R(q^2, \nu)} \right]$$

where E , p , E' , and p' are the incident and scattered muon laboratory energies and momenta, $q^2 = 2(EE' - pp' \cos\theta - m_\mu^2)$, $\nu = E - E'$, θ is the muon scattering angle, m_μ the muon mass, R the ratio of the total cross section on protons of scalar and transverse virtual photons, and α is the fine-structure constant.

The value of R has been determined by comparing cross sections at the same q^2 and ν using the 219-GeV data and all earlier measurements at 97 and 147 GeV. First, the two sets of measurements made under different conditions at 147 GeV were combined. Then the three different energy measurements were compared and χ^2 minimized as a function of F_2 and R where an overlap occurred. Table I shows the best-fit values of R as a function of $x = 1/\omega$ where $x = q^2/2m_p\nu$ and m_p is the proton mass. The χ^2 distributions of smaller bins were summed and minimized to obtain an average value of R ; this average value

is 0.44, and the statistical error is ± 0.16 from a $\chi^2 = 159$ for 148 degrees of freedom. The systematic errors in the measurement of R are estimated to be ± 0.05 from the normalization, ± 0.10 from uncertainty in the acceptance, and ± 0.15 from uncertainty in track-finding efficiency. These errors have been combined in quadrature

TABLE I. Values of $R = \sigma_S/\sigma_T$ for bin-centered values of x . The errors are statistical. The systematic error is ± 0.19 .

x	$q^2 \text{ (GeV}^2\text{)}$	R
0.069	2-12.5	$-0.11^{+0.29}_{-0.20}$
0.025	2-6	$0.56^{+0.31}_{-0.23}$
0.009	1-3	$0.44^{+0.27}_{-0.22}$

with the statistical error to give the final result $R = 0.44 \pm 0.25$ for $0.009 < x < 0.1$ and $1 < q^2 < 12 \text{ GeV}^2$.

Since there are uncertainties in the value of R measured at low energy³ for $x > 0.1$ and since our data are insensitive to these uncertainties for $x > 0.1$ (as shown in Fig. 2), we use our value to derive our best estimate for $\nu W_2(q^2, \nu) \equiv F_2(x, q^2)$. Figure 1 shows the values of the proton structure function as a function of q^2 for various values of ω . The values are weighted to give correct values at the bin centers. Only statistical errors are shown. The overall systematic error is estimated to be 5%. There is agreement between these new measurements and our previous values which are also shown recalculated for $R = 0.44$. A consistency check shows that the data have a $\chi^2 = 148$ for 126 degrees of freedom. In Fig. 2 we have combined all the data for muon scattering on hydrogen to show the variation of F_2 with x for various q^2 ranges. The Massachusetts Institute of Technology-Stanford Linear Accelerator Center (MIT-SLAC) data^{4,5} are also shown. The sensitivity of our results at each point to a variation of R between 0.19 and 0.69 is also indicated. These results exhibit a now well-known pattern of scaling violation^{2,5,7} which may be




FIG. 1. F_2 for hydrogen as a function of q^2 for various ω bins for $R = 0.44$ showing the 219-GeV data (closed circles), the earlier Fermilab data (Ref. 2) recalculated for $R = 0.44$ (open circles), and the MIT-SLAC data (Ref. 4) (open triangles). Note the difference in scales and the suppressed zeroes for the ordinate. The solid lines are the fits with Eq. (1) for all Fermilab data.




FIG. 2. $F_2(x, q^2)$ for hydrogen as a function of x for various q^2 bins for $R = 0.44$ using all Fermilab data (open circles) and the MIT-SLAC data (open triangles, Ref. 4; crosses, Ref. 5). The upper and lower horizontal bars indicate the change in F_2 for $R = 0.69$ and 0.19 . The errors shown are statistical only.

characterized by a power-law dependence of F_2 on q^2 for various x values where

$$F_2(x, q^2) = F_2(x, q_0^2) (q^2/q_0^2)^b \tag{1}$$




FIG. 3. The scaling violation parameter b of Eq. (1) as a function of x for all Fermilab data (closed circles) and for the MIT-SLAC data (Ref. 4) (open circles). The upper and lower horizontal bars indicate the change in b for $R = 0.69$ and 0.19 . The errors are statistical only.

Figure 3 shows the variation of b as a function of x for the combined data for various assumed values of R for $q^2 > 1.0 \text{ GeV}^2$. The rise of F_2 with increasing q^2 at small x may be reduced with a smaller value of R but will not disappear for any reasonable variation of R as a function x and q^2 .

We have attempted to fit the combined Fermilab data to a global scale-noninvariant structure function suggested by one of us,⁸

$$F_2^p(x, q^2) = A(2 + \epsilon_0)x(1-x)^{1+\epsilon_0} + B \frac{14(4 + \epsilon_1)}{9(5 + \epsilon_1)} (1-x)^{3+\epsilon_1} \frac{q^2}{q^2 + m_0^2}, \quad (2)$$

where $\epsilon \equiv \kappa \ln[(q^2 + m_0^2)/m_0^2]$, $\epsilon_0 \equiv G_{00} + \epsilon$, $\epsilon_1 \equiv G_{10} + \epsilon$.

The functional form chosen is suggested by the simple phase-space approach of Bjorken⁹ combined with an explicit q^2 -dependent factor for the approach to scaling in the second term from Devenish and Schildknecht.¹⁰ The first term can be loosely associated with a bare photon interaction with a three-valence-quark (plus gluons) configuration. The second term is a diffractive component in which a quark-antiquark sea (plus gluons) is added to the valence quarks in the generalized-vector-dominance spirit. These forms were first suggested empirically by the scale-noninvariant properties of the data, but we note that our formula bears a resemblance to the one arrived at by Buras and Gaemers who have parametrized the results of asymptotically free gauge-theory predictions.¹¹ In particular, we differ principally in our parameter ϵ which we find varying approximately as the cube root of a logarithm, while Buras suggests a log-log behavior. The values of the parameters determined from a fit to the data are given in Table II.

Our new data enable us to make an improved calculation of the integral of the structure func-

TABLE II. Values of parameters for global fit [Eq. (2)] to $F_2^p(x, q^2)$ for all Fermilab data.

Parameter	$q^2 > 2.0 \text{ GeV}^2$	$q^2 > 0.2 \text{ GeV}^2$
A	0.656 ± 0.039	0.661 ± 0.035
B	0.395 ± 0.021	0.357 ± 0.009
κ	0.27 ± 0.20	0.33 ± 0.17
G_{00}	2.00 ± 0.72	1.69 ± 0.68
G_{10}	7.7 ± 1.9	7.2 ± 1.7
$m_0^2 \text{ (GeV}^2\text{)}$	1.02 ± 0.22	0.573 ± 0.052
χ^2/NDF	46/38	96/64

tion $\int F_2^p(x, q^2) dx$ and the equivalent Nachtmann moment¹² ($N=2$) in different q^2 bands. The procedure used has been modified from that described in Ref. 2 by a numerical integration of the MIT-SLAC data including the resonance contributions. The results are given in Table III which replaces Table III of Ref. 2, and are given with and without the elastic scattering contribution. The uncertainty in R is included in the systematic error. These integrals show a slight decrease with increasing q^2 .

The results of this work reinforce the conclusions suggested by the earlier muon-scattering data that $F_2(x, q^2)$ increases as x goes to zero for fixed q^2 greater than 2 GeV^2 and increases with q^2 for fixed x less than 0.1. This behavior combined with the decrease of F_2 with increasing q^2 for x greater than 0.2 is consistent with that predicted by quantum chromodynamics for a scale parameter $\Lambda = 500-700 \text{ MeV}$.¹³ The integrals of F_2^p show that the quarks carry about half of the total energy-momentum of the proton, the fraction falling slightly with q^2 .¹³ Unfortunately the present statistical and systematic uncertainties prevent any meaningful comparison of R with quantum-chromodynamic predictions.¹⁴

TABLE III. Values of the energy-momentum sum rule.

q^2 (GeV^2)	x_{\min}	$\int_{x_{\min}}^{0.25} F_2^p dx$ Fermilab data only	$\int_0^1 F_2^p dx$		$N=2$ Nachtmann moment		Errors for integrals	
			No elastic contribution	Including elastic contribution	No elastic contribution	Including elastic contribution	Statistical	Systematic
1-2	0.0025	0.0673 ± 0.0020	0.1876	0.2150	0.1831	0.2095	± 0.0004	± 0.0060
2-4	0.0042	0.0840 ± 0.0018	0.1892	0.1952	0.1870	0.1919	± 0.0008	± 0.0070
4-8	0.0083	0.0838 ± 0.0014	0.1861	0.1868	0.1843	0.1849	± 0.0016	± 0.0070
8-15	0.0222	0.0815 ± 0.0018	0.1716	0.1717	0.1708	0.1709	± 0.0030	± 0.0070
15-30	0.0500	0.0768 ± 0.0027	0.1681	0.1682	0.1676	0.1676	± 0.0049	± 0.0090
30-50	0.1111	0.0512 ± 0.0050	0.1661	0.1661	0.1657	0.1657	± 0.0149	± 0.0140

We would like to thank Richard Heisterberg in particular and the staffs of Fermilab, the Rutherford Laboratory, and our respective laboratories for their help in this experiment. This work was supported by the National Science Foundation under Contract No. MPS71-03-186, PHY-76-23245, by the U. S. Department of Energy under Contracts No. EY-76-C-02-3064, No. E(11-1)-1195, No. E-(40-1)-2504, and by the Science Research Council (United Kingdom).

¹H. L. Anderson *et al.*, Phys. Rev. Lett. **37**, 4, 1034(E) (1976).

²H. L. Anderson *et al.*, Phys. Rev. Lett. **38**, 1456 (1977).

³L. N. Hand, in *Proceedings of the International Symposium on Lepton and Photon Interactions at High Energies, Hamburg, 1977*, edited by F. Gutbrod (DESY, Hamburg, Germany), p. 417.

⁴E. M. Riordan *et al.*, SLAC Report No. SLAC-PUB-1634, 1975 (unpublished).

⁵R. E. Taylor and W. B. Atwood kindly made available

a complete set of the electron-scattering results from the MIT-SLAC experiments.

⁶C. Chang *et al.*, Phys. Rev. Lett. **35**, 901 (1975).

⁷K. W. Chen, in *Proceedings of the International Symposium on Lepton and Photon Interactions at High Energies, Hamburg, 1977*, edited by F. Gutbrod (DESY, Hamburg, Germany), p. 467.

⁸T. B. W. Kirk, Fermilab Report No. TM-791, 1973 (unpublished).

⁹J. D. Bjorken and E. A. Paschos, Phys. Rev. **185**, 1975 (1969).

¹⁰R. Devenish and D. Schildknecht, Phys. Rev. D **14**, 93 (1976).

¹¹A. J. Buras and K. J. F. Gaemers, Nucl. Phys. **B122**, 249 (1978).

¹²O. Nachtmann, Nucl. Phys. **B63**, 237 (1973); and **B73**, 455 (1974).

¹³G. C. Fox, Nucl. Phys. **B131**, 197 (1977); H. I. Anderson, H. S. Matis, and L. C. Myriantopoulos, Phys. Rev. Lett. **40**, 1961 (1978).

¹⁴J. Hinchliffe and C. H. Llewellyn Smith, Nucl. Phys. **B128**, 93 (1977); A. J. Buras, E. G. Floratos, D. A. Ross, and C. T. Sachrajda, Nucl. Phys. **B131**, 308 (1977).